

Sweetpotato

Introduction

The terms “sweetpotato” and “yam” are often used interchangeably; however, they are actually two entirely different crops. Only sweetpotatoes (*Ipomoea batatas*) are grown in the U.S.; yams (*Dioscorea* spp.) are grown in the Caribbean and other tropical areas.

Marketing

The most profitable marketing opportunities for sweetpotatoes in Kentucky are through local fresh markets, such as farmers markets and on-farm stands. “U-Dig” sweetpotato sales, similar to U-Pick, are also possible in some areas. Currently (2010) there are about 200 acres of commercial sweetpotato production in the state. While sweetpotato processing has grown nationally in recent years and is dominated by large processors, there are no significant processing markets available in Kentucky.

Market Outlook

U.S. sweetpotato use per capita has increased significantly during the last decade and was estimated at 5.7 pounds in 2010. This represented a 35% increase over 2000 levels. Several factors, including increases in consumption of sweetpotato fries, have contributed to this increase. Sweetpotatoes regained some popularity with their emphasis as a “lower-carbohydrate potato” in the early 2000s; however, high antioxidant levels in sweetpotato skins have contributed to consumption staying strong after the low-



carb diet craze. Sweetpotato consumption is highest among Americans over 60, and sweetpotatoes may have special appeal to aging, health-conscious baby boomers. White and purple flesh sweetpotatoes, as well as organic sweetpotatoes, are possible market niches that Kentucky growers could fill. Higher market prices occur during the winter months so growers who are able to store their crop will be able to increase profits substantially.

Production Considerations

Cultivar selection

There are hundreds of sweetpotato cultivars available, including heirloom varieties. Sweetpotatoes differ in such horticultural characteristics as root shape, skin color (copper, rose, red, white, or purple), and flesh color (orange, deep orange, purple, or cream). Flavor (sweetness) and flesh moisture can also vary. Dry flesh types are often preferred by Asian and Hispanic consumers. Because some cultivars require the lengthy growing season found farther south, it is important to select only cultivars known to be well-adapted for the Kentucky

**Crop Diversification
& Biofuel Research
Education Center**

growing season. Resistance to various diseases (e.g. root knot nematode, internal cork, and/or Fusarium wilt) and insects (e.g. wireworm) is available in some cultivars. Commercial growers should choose varieties with the qualities in demand by the intended market.

Site selection and planting

Sweetpotatoes grow best on medium to light sandy soils that are well drained. While this crop can be grown in heavier soils, roots may be rough-skinned and irregular as a result. Fields high in organic matter should be avoided. Sweetpotatoes should not be grown on the same land more often than once every 3 years. Avoid fields with a history of difficult to control perennial weeds.

Propagation is from vine cuttings, which are referred to as “slips.” While certified disease-free slips can be purchased, it is often more economical for growers to propagate their own stock. Ten to 12 bushels of disease-free sweetpotatoes should be bedded to produce enough slips for one acre. Ordinarily 16 to 20 square feet of bed surface will be needed for each bushel. Sweetpotatoes are usually bedded about 7 weeks before field-setting time.

Sweetpotatoes are cold-sensitive and are planted after all danger of frost has passed. Slips may be transplanted to the field by hand, but many Kentucky growers use a one-row tobacco setter. About 15,000 transplants (slips) are set per acre. Providing an even supply of water during the first 40 days after planting is especially important for quality root development. An uneven water supply can result in growth cracks; drought conditions may reduce yields; and excess moisture may injure roots. Additionally, watering during this critical period can help plants survive later water-related stresses.

Pest management

The main insect pests are those that feed on the roots, such as wireworms, flea beetle larvae, and sweetpotato weevils. Diseases include black rot and scurf, Fusarium wilt, root knot nematodes,

and post-harvest rots. Resistant cultivars, crop rotations, sanitation, and weed management are important tools in disease and insect management. Pesticide applications may be necessary in some seasons.

Harvest and storage

Sweetpotatoes should be harvested when sufficient 6- to 8-ounce potatoes are found in the hill. A good practice is to clip the vines before harvesting so they do not get in the way during harvest, resulting in less damage to the potatoes. A turn plow or a potato plow can be used to expose the roots with the least possible injury. Potatoes are graded in the field and then placed in containers that are to be put into storage. For large scale production, mechanical harvesting machinery must be used.

Following harvest, sweetpotatoes need to undergo a curing process to promote the healing of wounds (for example, from digging). Curing protects roots from many storage diseases and increases the post-storage lifetime of the root. Additionally, curing improves root flavor and texture. Curing is best accomplished at a temperature of 85° F and relative humidity of 80% to 90%. After curing, sweetpotatoes may be stored for 4 to 7 months under the proper conditions, including ventilation. Sweetpotatoes are cleaned, either by brushing or washing, and then sometimes waxed before packing into boxes, crates or baskets for market.

Labor requirements

Labor needs per acre are approximately 16 hours for production, 40 hours for harvest, and 40 to 50 hours for post-harvest handling.

Economic Considerations

Initial investments include land preparation and the purchase of “slips” or sweetpotatoes for propagation. Total preharvest expenses are approximately \$1,600 per acre. Harvest and marketing costs for a 425-box yield will add approximately \$2,000 to variable costs. Total costs, including variable and fixed, are estimated

just over \$4,000 per acre. Since returns vary depending on actual yields and market prices, the following per acre returns to land and management are based on three different economic scenarios.

Conservative estimates represent the University of Kentucky's average cost and return estimates for wholesale production in 2010. Wholesale sweetpotato prices were stronger in 2010 than some past years; these returns are in line with those experienced by a mixture of direct and wholesale marketing in other recent years.

PESSIMISTIC	CONSERVATIVE	OPTIMISTIC
\$740	\$2,895	\$3,775

Selected Resources

- Sweetpotato Production for Kentucky, ID-195 (University of Kentucky, 2012) *1 MB file*
<http://www.ca.uky.edu/agc/pubs/id/id195/id195.pdf>
- Vegetable and Melon Budgets (University of Kentucky, 2013)
<http://www.uky.edu/Ag/cdbrec/vegbudgets13.html>
- Vegetable Production Guide for Commercial Growers, ID-36 (University of Kentucky)
<http://www.ca.uky.edu/agc/pubs/id/id36/id36.htm>

- Commercial Sweet Potato Production in Mississippi (Mississippi State University, 2002)
<http://msucare.com/pubs/publications/p1678.html>
- Estimated Costs and Returns for Sweet Potatoes (Fresh market, Irrigated) (Clemson University, 2009)
<http://cherokee.agecon.clemson.edu/swtpot6.pdf>
- Growing Sweet Potatoes Commercially (North Carolina Sweet Potato Commission)
<http://www.ncsweetpotatoes.com/sweet-potato-industry/growing-sweet-potatoes-in-north-carolina/>
- Guide to Sweetpotato Production in Alabama ANR-982 (Alabama Cooperative Extension, 2006) *4.96 MB file*
<http://www.aces.edu/pubs/docs/A/ANR-0982/ANR-0982.pdf>
- Harvesting and Curing Sweetpotatoes ANR-1111 (Alabama Cooperative Extension, 2004)
<http://www.aces.edu/pubs/docs/A/ANR-1111/ANR-1111.pdf>
- Sweet Potato Varieties (North Carolina Sweet Potato Commission)
<http://www.ncsweetpotatoes.com/sweet-potatoes-101/sweet-potato-varieties/>
- Sweetpotato: Organic Production (ATTRA, 2005)
<https://attra.ncat.org/attra-pub/summaries/summary.php?pub=32>

Reviewed by John Strang, Extension Specialist (Issued 2002, Revised 2006)
Reviewed by Shawn Wright, Extension Specialist (Revised December 2010)
Photo by Charles Averre, North Carolina State University, Bugwood.org

December 2010

For additional information, contact your local [County Extension](#) agent